

Lensing and the luminosity of Gamma-ray bursts and their hosts

C.A. Scharf¹ and K.C. Sahu²

Space Telescope Science Institute, 3700 San Martin Drive, Baltimore MD 21218, USA

Received –; accepted –

¹scharf@stsci.edu

²ksahu@stsci.edu

ABSTRACT

The quoted intrinsic luminosity of objects has a dependency on the assumed cosmology, and the (less often considered) assumed gravitational lensing due to the matter content of the viewing beam; the so called ‘filled’ or ‘empty’ beam cases.

We consider the implications of filled vs empty beam assumptions for the derived total luminosities of recent gamma-ray bursts (GRBs) in which counterparts at cosmological distances have been confirmed. Conversion factors between filled and empty beam bolometric luminosities are presented graphically for a range of cosmologies and redshifts. The tendency for sources to be most probably demagnified further supports the need for non-isotropic GRB emission in neutron star merger models, or the need for at least one massive star in other models. In the most extreme case the true energy of GRB971214 at $z = 3.418$ could be as high as $\sim 8 \times 10^{53}$ ergs, a factor ~ 2 more than previously estimated. Similarly, the effect may account for some of the observed bias towards fainter host magnitudes.

Subject headings: gamma rays:bursts — cosmology: gravitational lensing

1. Introduction

The intrinsic luminosity of an object is estimated using the luminosity distance to the object. This distance is usually derived assuming a Robertson-Walker model of the Universe where the matter is uniformly and homogeneously distributed in space, the so-called filled-beam distance ($d_{L-filled}$). The matter, as we know it however, is comprised of compact objects such as stars and galaxies, and has a clearly inhomogeneous distribution. The opposite extreme to the filled-beam case is the so called empty-beam case, where all the matter is in the form of compact objects far from the the line-of-sight; the derived distance being called the empty-beam distance ($d_{L-empty}$). In this latter case the weak lensing of the rest of the universe de-magnifies the source relative to the filled-beam case. Although reality may be somewhere between these two extremes (simulations indeed show that the intermediate case is more representative of the actual matter distribution, see below) it is clearly important to see to what extremes the matter distribution can affect the derived luminosity.

Recently several authors have re-examined this phenomenon with detailed investigations of the expected range of magnification effects expected in various model cosmologies (e.g. Wambsganss, Cen & Ostriker 1997, Kantowski 1998, Holz 1998). These results, which build on earlier works such as Ostriker & Vietri (1986), and Dyer & Roeder (1973), demonstrate that in most commonly acceptable cosmologies the probability distribution of the observed magnitude is strongly peaked intermediate to the empty and filled-beam cases (the skewness depending largely on whether the lensing matter is distributed as points or more diffusely, the former skewing magnitudes significantly towards the empty-beam, de-magnified, case). However a tail to high magnification exists, so sources may also be brightened.

In this short paper we point out the relevance of this effect to GRBs and their host galaxies, we also present a simple diagram for reading off the extreme case of empty vs filled

beam demagnification factors for various cosmologies. We present the estimated energy output of several GRBs and the range allowed due to de-magnification effects, as well as the range of effects on the host galaxy magnitudes.

2. Empty vs filled luminosity factors

Since the bolometric luminosity for a given flux/apparent magnitude varies as the square of the luminosity distance (d_L) we estimate here the maximum possible difference in inferred luminosity as the ratio $d_{L-empty}^2/d_{L-filled}^2$. In figures 1 and 2 we plot this luminosity factor as a function of the cosmological density parameter, Ω_0 , and redshift, z , for both open and flat (non-zero cosmological constant, $\lambda_0 \neq 0$) universes. The relevant equations are given in Dyer & Roeder (1973) and Fukugita et al (1992).

Several features are worth noting. First, as expected, the higher the matter density the larger the luminosity factor between empty and filled beam estimates since the empty beam will be in higher contrast to the rest of the universe as Ω_0 increases. Second, the luminosity factors are fairly small for $z \lesssim 1$, at most $\sim 20\%$, but by $z \sim 4$ can rise as high as $\sim 300\%$ (for $\Omega = 1$).

3. Implications for GRBs

In Table 1 we summarize the current data on GRBs with identified hosts and measured redshifts (note: GRB 970228 has a redshift estimated from photometric colours; Sahu et al. 1997a, Fruchter et al., 1998). In the final column we give the range of intrinsic GRB energies if the sources have been demagnified. These energies represent the extreme case of empty-beam demagnification for flat ($\Omega_0 + \lambda_0 = 1$) cosmologies with $\Omega_0 = 0.2$ or $\Omega_0 = 1$. It should be noted that the $\Omega = 1$ values are given to illustrate the effect of the

overall cosmology. The burst GRB971214 is subject to the largest possible underestimate of intrinsic energy (Odewahn et al. 1998), by a factor of 2, to a maximum of $\sim 8 \times 10^{53}$ ergs.

4. Host galaxy magnitude distribution

Recently Hogg & Fruchter (1998) have shown how the distribution of GRB host galaxy magnitudes ($24.4 \leq R \leq 25.8$, and one with $R = 14.3$) appears to be biased towards the fainter end of the expected magnitude distribution, based on measured and inferred redshifts. The bias results in an apparent deficit in hosts with $21 \leq R \lesssim 24$, albeit at low significance. While there are clearly other potential reasons for such a bias the possibility that at least some fraction of these high- z objects have been de-magnified to lower apparent magnitudes cannot be discounted. A luminosity factor of ~ 1.5 (a plausible ‘typical’ maximum factor over the redshift range $1 < z < 3$, a range expected from the star-formation history of the Universe (Sahu et al. 1997b; Hogg & Fruchter, 1998)) results in a magnitude shift of ~ -0.44 which could account for a significant amount of the faint end bias. Even a more realistic factor of 1.2 (-0.2 magnitudes) which might be expected from detailed simulation, e.g. Wambsganss, Cen, & Ostriker (1998) helps alleviate the dearth of brighter hosts.

5. Discussion

Since GRB 971214 is the only GRB with an observed redshift > 2 , this deserves special mention. If the luminosity distance for high- z GRBs is more accurately described by the empty-beam case, then the quoted energy of GRB971214 could be higher by a factor ~ 2 ($\Omega = 0.2$), implying an energy output of some $\sim 8 \times 10^{53}$ ergs. If the energy is isotropic, the released energy is too high for most models, including the neutron star merger models.

This would indicate that the GRB emission may be strongly beamed, which would be a natural consequence of relativistic expansion, or may require a neutron star-massive black hole merger model, or a hypernova model for the GRB. Furthermore, GRB host galaxies to the fainter side of the expected magnitude distribution could easily be explained by such a variation in the luminosity distance.

Detailed studies of cosmological lensing indicate that the most probable effect is a milder demagnification of high- z sources than these extreme values, with a probability tail of positive magnification leading up to strong lensing events. Given the extreme nature of GRBs however, where the observed energies are already too large compared to the predictions from many models, the possibility that we are systematically underestimating the intrinsic energy release is perhaps the more relevant issue. Filled beam estimates of intrinsic luminosity should therefore be considered subject to an unknown amount of scatter, tending towards underestimates of the true value by as much as the factors presented here.

M. Donahue is thanked for useful discussions and C.A.S. acknowledges support through NASA grant NAG5-3257

REFERENCES

- Bloom, J.S. et al 1998, preprint (astro-ph/9808319)
- Bloom, J.S. et al 1998, preprint (astro-ph/9807050)
- Dyer, C. C., Roeder, R. C. 1973, ApJ, 180, 31
- Fruchter, A.S., et al., 1998, ApJ, submitted (astro-ph/9807295)
- Fukugita, M., Futanase, T., Kasai, M., Turner, E.L. 1992, ApJ, 393, 3
- Hogg, D.W., Fruchter, A.S. 1998 ApJ, submitted (astro-ph/9807262)
- Holz, D.E. 1998, Preprint (astro-ph/9806124)
- Kantowski, R. 1998, Preprint (astro-ph/9802208)
- Kulkarni, S.R. et al 1998, Nature, 393, 35
- Metzger, R.M. et al 1997, Nature, 387, 878
- Odewahn, S.C., et al., 1998, ApJ, submitted (astro-ph/9807212)
- Ostriker, J.P., Vietri, M. 1986, ApJ, 300, 68
- van Paradijs, J. et al 1997, Nature, 386, 686
- Sahu, K.C., et al., 1997a, Nature, 387, 476
- Sahu, K.C., et al., 1997b, ApJ, 489, L127
- Wambsganss, J., Cen, R., Ostriker, J.P. 1998, ApJ, 498, 29

Fig. 1.— Contour map of the luminosity factor between empty and filled beam models for open universes ($\lambda_0 = 0$) as a function of redshift. The contours are at constant $d_{L-empty}^2/d_{L-filled}^2$ and are labeled. Dashed contours run from factors 1.1, 1.2, 1.3 to 1.4, solid contours run from 1.5 in steps of 0.5

Fig. 2.— Contour map of the luminosity factor between empty and filled beam models for flat universes ($\Omega_0 + \lambda_0 = 1$) as a function of redshift. Contours are as for Figure 1

Name	Redshift	Energy (ergs) ^a	Maximum energy ‘empty-beam’
		($\Omega_0 = 1$), ($\Omega_0 = 0.2$, $\lambda_0 = 0.8$)	($\Omega_0 = 1$), ($\Omega_0 = 0.2$, $\lambda_0 = 0.8$)
980703 ^b	0.966	$0.7 - 1.4 \times 10^{53}$	$0.9 - 1.5 \times 10^{53}$
971214 ^c	3.418	$1.44 - 4 \times 10^{53}$	$3.8 - 8 \times 10^{53}$
970508 ^d	0.835	$5.18 - 9.1 \times 10^{51}$	$6.22 - 10 \times 10^{51}$
970228 ^e	≥ 1	$\geq 0.69 - 1.4 \times 10^{51}$	$0.9 - 1.54 \times 10^{51}$
980425 ^f	0.0085	8.1×10^{47}	8.1×10^{47}

^aDerived energies assuming $H_0 = 65 \text{ kms}^{-1}$

^bBloom et al 1998a

^cKulkarni et al 1998

^dMetzger et al 1997

^evan Paradijs et al 1997

^fBloom et al 1998b

Note. — GRBs with identified hosts and measured redshifts. Full and empty beam energies are estimated for $\Omega_0 = 1$ and $\Omega_0 = 0.2$, $\lambda_0 = 0.8$ cosmologies.



